

The Geometry of Radiative Transfer

Christian Lessig,^{1,2} Alex L. Castro,³ and Eugene Fiume²

¹*Computing $\times$ Mathematical Sciences, Caltech, Pasadena, CA, USA*

²*Dynamics Graphics Project, University of Toronto, ON, Canada*

³*Pontifícia Universidade Católica do Rio de Janeiro, Rio de Janeiro, Brazil*

(Dated: March 8, 2013)

We present the geometric structure of radiative transfer and the symmetries associated with its known conserved quantities. Our geometrization exploits recent work in the literature that enables the Hamiltonian formulation of radiative transfer to be obtained from a phase space representation of electromagnetic theory. Cosphere bundle reduction yields the traditional description over the space of “positions and directions”, and geometrical optics arises as a special case when energy is disregarded. It is also shown that, in idealized environments, radiative transfer is a Lie-Poisson system with the group of canonical transformations as configuration space and symmetry group.

PACS numbers: 42.15.—i, 46.05.+b, 47.10.—g

Radiative transfer describes the transport of electromagnetic energy in macroscopic environments when polarization effects are neglected [1]. The theory originates in work by Bouguer [2, 3] and Lambert [4] in the 17th century, where the measurement of light intensity was first studied systematically, and was later extended, for example to include transport effects [5–8]. Despite these efforts, however, to this day radiative transfer remains phenomenological, largely without connection to the fundamental theories of light, and without a suitable modern mathematical formulation—despite its importance in a multitude of fields, such as medical imaging, remote sensing, computer graphics, atmospheric science, climate modelling, and astrophysics.

In this letter, we explain the physical foundations of radiative transfer in media with varying refractive index and we study its Hamiltonian structure and symmetries using a modern mathematical formulation, see Fig. 1. Exploiting recent work in the literature, we derive radiative transfer from electromagnetic theory using an ansatz that resembles semi-classical analysis. The Wigner transform is employed to lift Maxwell’s equations from configuration space $Q \subset \mathbb{R}^3$ to phase space T^*Q . Restricting the dynamics on T^*Q to a non-zero energy level, one obtains at the short wavelength limit a transport equation for polarized light, and further neglecting polarization leads to radiative transfer theory in a Hamiltonian formulation. This derivation uncovers the phase space light energy density $\ell \in \text{Den}(T^*Q)$ as central quantity of radiative transfer theory, and shows that radiance, which classically plays this role, is meaningful only in the context of measurements, in the setting Lambert considered when he introduced the concept. With the Hamiltonian formulation of radiative transfer on 6-dimensional phase space T^*Q , the classical 5-dimensional description is obtained by reduction to the cosphere bundle $S^*Q = (T^*Q \setminus \{0\})/\mathbb{R}^+$ where time evolution is described by contact dynamics. The conserved quantity associated with this symmetry is the light frequency, un-

covering the structure of a long known conservation law. Fermat’s principle is obtained from the Hamiltonian formulation of radiative transfer through a non-canonical Legendre transform. Geometrical optics hence arises as a special case of the theory. When for a fixed environment Q the light energy density $\ell_t \in \text{Den}(T^*Q)$ at time t is considered as a configuration of radiative transfer, and when one assumes that the Hamiltonian flow of the system is defined globally, the configuration space of radiative transfer becomes the group $\text{Diff}_{\text{can}}(T^*Q)$ of canonical transformation. This makes ideal light transport a Lie-Poisson system, and reveals a remarkable similarity to ideal fluid dynamics, cf. Table I. The symmetry associated with the Lie-Poisson structure is the conservation of light energy density along trajectories in phase space, providing a modern rationale for the classical law of “conservation of radiance along a ray” whose symmetry was previously not known. The geometrization and physical foundations of radiative transfer presented in this letter embed the theory into modern physics and they enable—and warrant—a fresh look at new and old questions.

The physical foundation of radiative transfer theory is provided by the macroscopic Maxwell’s equations, Hamilton’s equations for electromagnetic field theory [9, p. 24], which in a source free region $Q \subset \mathbb{R}^3$ are given by

$$\frac{\partial}{\partial t} \begin{pmatrix} \vec{E} \\ \vec{H} \end{pmatrix} = \begin{pmatrix} 0 & -\frac{1}{\epsilon} \nabla \times \\ \frac{1}{\mu} \nabla \times & 0 \end{pmatrix} \begin{pmatrix} \vec{E} \\ \vec{H} \end{pmatrix} \quad (1)$$

where $\epsilon : Q \rightarrow \mathbb{R}$ and $\mu : Q \rightarrow \mathbb{R}$ are the electric permittivity and magnetic permeability, respectively. With $U = (\vec{E}, \vec{H})^T$, Eq. 1 can be written as

$$\dot{U}^\epsilon = P^\epsilon U^\epsilon \quad (2)$$

where P^ϵ is the Maxwell operator. In Eq. 2 we also introduced the scale parameter $\epsilon = \lambda/d_n$ with λ being the wavelength and d_n the average distance over which the refractive index $n = \sqrt{\epsilon\mu} : Q \rightarrow \mathbb{R}$ varies. Macroscopic environments are characterized by $\lambda \ll d_n$, which warrants considering the short wavelength limit $\epsilon \rightarrow 0$. The




FIG. 1. The physical foundations and geometric structure of radiative transfer theory.

observable of interest for radiative transfer theory is the electromagnetic energy density

$$\mathcal{E}^\varepsilon(q, t) = \|U_{\varepsilon, \mu}^\varepsilon\|^2 = \frac{\varepsilon}{2} \|\vec{E}^\varepsilon\|^2 + \frac{\mu}{2} \|\vec{H}^\varepsilon\|^2 \quad (3)$$

for $\varepsilon \rightarrow 0$ and when polarization is neglected. The classical approach to study short wavelength asymptotics is the Wenzel-Kramers-Brillouin (WKB) approximation. However, the ansatz yields well-defined solutions only until caustics form, when the approximation becomes multi-valued, and the initial conditions must satisfy the WKB form $u^\varepsilon(q, t) = a(q, t) e^{iS(q, t)/\varepsilon}$. As shown recently [10], these limitations can be circumvented by studying the short wavelength limit on phase space $T^*Q \cong \mathbb{R}^3 \times Q \subset \mathbb{R}^3$. The electromagnetic field U^ε can be lifted to T^*Q using the Wigner transform, yielding a 6×6 “matrix” density $W^\varepsilon = W[U^\varepsilon](q, p)$. Eq. 2 then becomes

$$\dot{W}^\varepsilon = -\{\{p^\varepsilon, W^\varepsilon\}\}_{\text{MB}} \quad (4)$$

where $p^\varepsilon = p_0 + \varepsilon p_1 + \varepsilon^2 p_2 + \dots$ is the symbol of the Maxwell operator P^ε , the dispersion matrix for the system [12, Eq. 3.63], and the bracket $\{\{, \}\}_{\text{MB}}$ on the right hand side is the matrix-valued Moyal bracket [15, Chapter 16.3]. In contrast to the scalar case where the Moyal bracket satisfies $\{\{, \}\}_{\text{MB}} \xrightarrow{\varepsilon \rightarrow 0} \{, \}$ and reduces to the Poisson bracket $\{, \}$ as $\varepsilon \rightarrow 0$, in the matrix valued case one has

$$\{\{p^\varepsilon, W^\varepsilon\}\} = \frac{1}{\varepsilon} (p^\varepsilon W^\varepsilon - W^\varepsilon p^\varepsilon) + \dots + O(\varepsilon) \quad (5)$$

with the first term diverging as $\varepsilon \rightarrow 0$ since matrix multiplication does not commute. For the short wavelength limit, this can be avoided by restricting the dynamics to an eigenspace, or allowed energy level, of the zeroth order Maxwell symbol p_0 , the analogue of the Bohr-Sommerfeld quantization condition [16]. The eigenvalues of p_0 are [12]

$$\lambda_0 = 0 \quad \lambda_1 = \frac{c}{n(q)} \|p\| \quad \lambda_2 = -\frac{c}{n(q)} \|p\| \quad (6)$$

with each having multiplicity two. Projection onto the eigenspace associated with a non-trivial eigenvalues $\lambda_a, a \in \{1, 2\}$ and taking the limit $\varepsilon \rightarrow 0$, one obtains [11]

$$\dot{W}_a^0 + \{\lambda_a, W_a^0\} = [W_a^0, F_a^0] \quad (7)$$

where the projection W_a^0 of the limit Wigner distribution W^0 onto the 2-dimensional eigenspace is given by

$$W_a^0 = \Pi_a W^0 \Pi_a = \frac{1}{2} \begin{bmatrix} I + Q & U + iV \\ U - iV & I - Q \end{bmatrix} dq \, dp. \quad (8)$$

W_a^0 is formed by the Stokes parameters I, Q, U, V for polarized light for the plane wave propagating in “direction” $p \in T_q^*Q$, and this also provides the physical interpretation of p as the wave vector $p = n(q)/c\omega \bar{s}$, with ω being the angular frequency and $\bar{s}$ a unit vector. With Eq. 8, the right hand side of Eq. 7 can be interpreted as describing the change in polarization during the transport by the Poisson bracket $\{\lambda_a, W_a^0\}$. Radiative transfer does not consider polarization. With the Stokes parameters for unpolarized light, $Q, U, V = 0$, Eq. 7 becomes

$$\dot{\ell} = -\{\ell, H\}. \quad (9)$$

The Hamiltonian $H \in \mathcal{F}(T^*Q)$ in Eq. 9 is given by the eigenvalue λ_a , that is

$$H(q, p) = \pm \frac{c}{n(q)} \|p\| \quad (10)$$

so that the associated Hamiltonian flow is determined by the refractive index $n(q)$. The trace $\ell = \text{tr}(W_i^0) = \mathcal{L}(q, p) \, dq \, dp \in \text{Den}(T^*Q)$ is related to the electromagnetic energy density $\mathcal{E}(q)$ by the fiber integral

$$\mathcal{E}(q) = \int_{T_q^*Q} \ell = \int_{T_q^*Q} \mathcal{L}(q, p) \, dp, \quad (11)$$

and the phase space density ℓ is hence the central quantity in radiative transfer theory, representing the energy of the electromagnetic field at the short wavelength

limit when polarization effects are neglected. We denote $\ell \in \text{Den}(T^*Q)$ as *phase space light energy density* or *light energy density*.

To relate $\ell \in \text{Den}(T^*Q)$ to radiance, the central quantity in the classical formulation of radiative transfer, we have to consider measurements, the question Lambert was studying when he introduced the concept [4]. Using standard results from tensor calculus, one obtains for the energy E flowing through a 2-dimensional surface $\mathcal{M}$ in a time interval $[t_1, t_2]$ that [17, Chapter 3.2.6]

$$E = \int_{t_1}^{t_2} \int_{T^-\mathcal{M}} i_{X_H} \ell \quad (12a)$$

$$= \int_{t_1}^{t_2} \int_{\mathcal{M}} \frac{c}{n(q)} \int_{T_q^-\mathcal{M}} \mathcal{L}(q, p) (\bar{p} \cdot \bar{n}) dA d\bar{p} dt \quad (12b)$$

where $\bar{p}$ is a unit vector, $\bar{n}$ the surface normal of $\mathcal{M}$, and $T_q^-\mathcal{M}$ the positive half-space of T_q^*Q as defined by the surface normal $\bar{n}|_q$. Parametrizing the light energy density in spherical coordinates [17, Chapter 3.2.5], infinitesimally the measurement can be written as

$$\mathcal{L}(q, \bar{p}, \nu) (\bar{p} \cdot \bar{n}) dA d\bar{p} = \bar{n} \cdot (\mathcal{L}(q, \bar{p}, \nu) \bar{p} dA d\bar{p} d\nu). \quad (13)$$

When no measurement surface, and hence no normal $\bar{n}$, is fixed an infinitesimal measurement is thus given by

$$\Lambda = \mathcal{L}(q, \bar{p}, \nu) dA_\perp d\bar{p} d\nu \quad (14)$$

where $dA_\perp(\bar{p}) = \bar{p} dA$ is the standard area form for a surface orthogonal to the flow direction $\bar{p}$. The differential 2-form $\Lambda \in \Omega^2(Q)$ provides a modern interpretation of classical radiance, with the cosine term $(\bar{p} \cdot \bar{n})$, prevalent in the classical literature, arising from the pullback of Λ onto a surface with normal $\bar{n}$.

Remark 1. It has often been overlooked that radiance is meaningful only in the context of measurements, which led to considerable confusion as in [18], while the quantity that is naturally transported in radiative transfer is the phase space light energy density $\ell \in \text{Den}(T^*Q)$.

Eq. 9 describes radiative transfer theory as a Hamiltonian system on 6-dimensional phase space T^*Q . In the literature, however, the theory is usually defined over the 5-dimensional space of “positions and directions”. The connection between the two descriptions can be obtained by considering the symmetry associated with the well known conservation of frequency. Since the momentum p is proportional to frequency ν , the associated Lie group action is $m_\alpha(q, p) = (q, \alpha p)$ for the multiplicative group $(\mathbb{R}^+, \cdot)$ acting on the fibers T_q^*Q . Moreover, since the Hamiltonian in Eq. 10 is homogeneous of degree one in the momentum, the dynamics can be reduced from T^*Q to the 5-dimensional cosphere bundle [19, App. 4]

$$S^*Q = (T^*Q \setminus \{0\}) / \mathbb{R}^+ \quad (15)$$

where the time evolution is a contact Hamiltonian flow along the Reeb vector field [20]. The cosphere bundle

$$\begin{array}{ccc} \hat{L} = n^2(q) & \xleftrightarrow[\text{transform}]{\text{Legendre}} & \hat{H} = g^{ij} \hat{p}_i \hat{p}_j \\ \uparrow & & \downarrow \\ L = n(q) & & \sqrt{\hat{H}} = \frac{c}{n(q)} \|\hat{p}\| \\ & & \downarrow \\ & & H = \omega \hat{H} = \frac{c}{n(q)} \|p\| \end{array}$$

FIG. 2. Non-canonical Legendre transform relating radiative transfer to geometrical optics given by Fermat’s principle.

S^*Q , which can be identified with the sphere bundle S^2Q using a metric, provides a modern interpretation for the classical space of “positions and directions”, and the Hamiltonian being homogeneous of degree one explains why a description on S^*Q is possible.

Remark 2. A question open to this date is the relationship between radiative transfer and geometrical optics. The connection can be established with the Legendre transform of Fermat’s principle, cf. Fig. 2. As is well known, directly performing the transform for the Lagrangian $L = n(q)$ leads to a vanishing Hamiltonian [21]. Following Arnold [19], we hence consider instead of length, given by $L = n(q)$, the geometrical energy of a light path, given by $\hat{L} = n^2(q)$. This Lagrangian can be interpreted as a diagonal metric $g_{ij} = n^2(q)/c^2 \delta_{ij}$ and the associated geodesic flow is equivalently described by the Hamiltonian $\hat{H} = g^{ij} \hat{p}_i \hat{p}_j$ [22, p. 51] where $\hat{p}$ is the canonical momentum which is related to the kinetic momentum by $p = \omega \hat{p}$. Reverting the transition from path length to path energy, and including the factor of ω corresponding to energy we obtain for the Hamiltonian again Eq. 10. If we trace the flow in Fig. 2 backwards, we see that, from the point of view of radiative transfer, geometrical optics is a special case when energy transport is disregarded.

A central result in classical radiative transfer theory is the “conservation of radiance along a ray” [23]. The symmetry associated with this conservation law becomes apparent when the light energy density ℓ_t at time t , for all $(q, p) \in T^*Q$, is considered as one configuration of the system. Time evolution can then be described by the pullback $\ell_t = \eta_t^* \ell_0$ by the time t flow map η_t along the Hamiltonian vector field X_H arising from Eq. 10. When X_H is defined globally, the set of all maps η_t forms the infinite dimensional Lie group $\text{Diff}_{\text{can}}(T^*Q)$ [24] of canonical transformations, and with respect to ℓ_0 all physically valid configurations ℓ_t can be described by an element η_t in the group. $\text{Diff}_{\text{can}}(T^*Q)$ can hence be considered as the configuration space of *ideal radiative transfer*, equipping the system with a Lie-Poisson structure, cf. Fig. 3. The Lie algebra $\mathfrak{g}$ of $\text{Diff}_{\text{can}}(T^*Q)$ can be identified with the generating Hamiltonian functions, $\mathfrak{g} \cong \mathcal{F}(T^*Q)$, in which case the dual Lie algebra $\mathfrak{g}^*$ becomes $\text{Den}(T^*Q)$ [26]. With $\mathfrak{g}^* \cong \text{Den}(T^*Q)$, it is natural to consider the light energy density ℓ as an element in




FIG. 3. The structure of Lie-Poisson systems. Classical examples of such systems are the rigid body, where the Lie group is $SO(3)$ (with a left invariant Hamiltonian), and the ideal Euler fluid, where the Lie-group is the group $\text{Diff}_\mu(Q)$ of volume preserving diffeomorphisms [24, 25].

$\mathfrak{g}^*$. Time evolution is then described by coadjoint action $\text{Diff}_{\text{can}}(T^*Q) \times \mathfrak{g}_+^* \rightarrow \mathfrak{g}_+^*$ in the Eulerian representation, which infinitesimally is [17, Sec. 3.3]

$$\dot{\ell} = \text{ad}_{\frac{\delta \mathcal{H}}{\delta \ell}}^* \ell = \text{ad}_H^* \ell = -\{\ell, H\}. \quad (16)$$

That the Poisson bracket describes infinitesimal coadjoint action $\text{ad}^* : \mathfrak{g} \times \mathfrak{g}^* \rightarrow \mathfrak{g}^*$ provides a rigorous justification for considering $\text{Diff}_{\text{can}}(T^*Q)$ as the configuration space for ideal radiative transfer [27, Sec. 6]. In Eq. 16, $\mathcal{H} \equiv \mathcal{H}[\ell]$ is the field Hamiltonian

$$\mathcal{H}[\ell] = \int_{T^*Q} \ell(q, p) H(q, p) dq dp \quad (17)$$

which is defined through the “single particle” Hamiltonian $H(q, p)$ in Eq. 10. With the light energy density as an element in the dual Lie algebra, it follows immediately from the general theory of Lie-Poisson systems that the momentum map $\mathbf{J}_R$ is the convective light energy density and that it is conserved [9, Theorem 11.4.1], cf. Fig. 3. By the change of variables theorem, this can be interpreted as conservation of light energy density along trajectories in phase space, providing a modern formulation and justification for “conservation of radiance along a ray”. Remarkably, with the Lie-Poisson structure a close analogy between ideal radiative transfer and ideal fluid dynamics exists, cf. Table I. Next to the flow on phase space T^*Q , one can obtain a functional analytic description, related to the operator formulation of radiative transfer [28], using Stone’s theorem, yielding $\ell_t = \eta_t^* \ell_0 = \mathcal{U}_t \ell_0$ where $\mathcal{U}_t$ is a unitary operator.

Our geometric formulation of radiative transfer and the identification of the Lie group structures underlying the known conservation laws clarifies and unifies earlier work in the literature. The use of tensor calculus overcomes the limitations of the current formulation, and also of earlier attempts that employed vector calculus [29] and measure theory [30] to obtain a modern mathematical foundation for radiative transfer. Our derivation of the Hamiltonian formulation of radiative transfer follows recent work in applied mathematics [11, 12] that employs

	fluid dynamics	radiative transfer
Lie group	$\text{Diff}_\mu(Q)$	$\text{Diff}_{\text{can}}(T^*Q)$
Lie algebra	$\mathfrak{X}_{\text{div}}(Q)$	$\mathfrak{X}_{\text{Ham}}(T^*Q)$
dual Lie algebra	$\omega \in \Omega^2(Q)$	$\ell \in \text{Den}(T^*Q)$
coadjoint action	$\dot{\omega} = -\mathcal{L}_v \omega$	$\dot{\ell} = -\mathcal{L}_{X_H} \ell$
conservation law	Kelvin’s theorem	conservation of ℓ

TABLE I. Correspondence between ideal fluid dynamics and ideal radiative transfer. The fluid velocity is denoted by $v \in \mathfrak{X}_{\text{div}}(Q)$ and $\omega \in \Omega^2(Q)$ is the fluid vorticity.

the Wigner transform to study short wavelength asymptotics, chiefly because of the simple and linear relationship in Eq. 11. The use of the Wigner transform parallels earlier approaches [31] [32]. The authors of [11, 12] put these on a rigorous mathematical basis, which is crucial to treat the delicate questions arising from the highly oscillatory nature of U^ε , and they elucidated many aspects that previously seemed serendipitous. Currently lacking is a treatment of the transition from Maxwell’s equations to radiative transfer from an algebraic and geometric point of view. Gérard and coworkers [11], for example, did not interpret their equations as matrix-valued Moyal bracket, and hence did not relate their results to the large body of work on quantization [35]. The status and shortcomings of many classical derivations of radiative transfer theory were recently summarized by Mishchenko [38].

An interesting extension of our geometric formulation would be the inclusion of dissipation and scattering. The classical Beer-Lambert law is naturally compatible with our description [17, Remark 3.29] but it would be interesting, and of considerable practical relevance, to extend this to more general scattering models, for example based on recent work on kinetic theory [39]. An important objective for future work should also be to develop the connection between our formulation of radiative transfer and work by Mishchenko [40, 41] where radiative transfer theory for media with scattering particles was derived from electromagnetic theory using an alternative approach.

We thank Mathieu Desbrun, Tudor Ratiu, Boris Khesin, and Jerry Marsden for discussions. C.L. and E.F. acknowledge support by the National Science and Engineering Council of Canada, as well as MITACS and GRAND National Centres of Excellence. C.L. was also supported by NSF grant CCF-1011944. A.C. thanks the University of Toronto and the Fields Institute for their hospitality and a postdoctoral fellowship during which this work was initiated.

-
- [1] G. C. Pomraning, *The Equations of Radiation Hydrodynamics* (Pergamon Press, 1973).
 - [2] P. Bouguer, *Essai d’optique sur la gradation de la lumière* (Claude Jombert, Paris, 1729).

- [3] P. Bouguer, *Traité d'Optique sur la gradation de la lumière* (H. L. Guerin et L. F. Delatour, Paris, 1760).
- [4] J. H. Lambert, *Photometry or On The Measure and Gradation of Light, Colors, and Shade*, edited by D. L. DiLaura (Illum. Eng. Soc., 2001).
- [5] E. Lommel, *Ann. der Phys.* **272**, 473 (1889).
- [6] O. D. Chwolson, *Bull. l'Acad. Imp. Sci. St. Pétersbourg* **33**, 221 (1889).
- [7] A. Schuster, *Astrophys. J.* **21**, 1 (1905).
- [8] K. Schwarzschild, *Nachr. Kgl. Ges. d. Wiss. zu Gött. Math. Phys. Klasse* **195**, 41 (1906).
- [9] J. E. Marsden and T. S. Ratiu, *Introduction to Mechanics and Symmetry*, 3rd ed. (Springer-Verlag, New York, 2004).
- [10] [11, 12], see also [13] and [14, Section 2,3] for an introduction.
- [11] P. Gérard, P. A. Markowich, N. J. Mauser, and F. Poupaud, *Comm. Pure Appl. Math.* **50**, 323 (1997).
- [12] L. Ryzhik, G. Papanicolaou, and J. B. Keller, *Wave Motion* **24**, 43 (1996).
- [13] C. Sparber, P. A. Markowich, and N. J. Mauser, *Asymptotic Anal.* **33**, 153 (2003).
- [14] S. Jin, P. Markowich, and C. Sparber, *Acta Numerica* **20**, 121 (2011).
- [15] H. Spohn, *Dynamics of Charged Particles and Their Radiation Field* (Cambridge University Press, 2004).
- [16] R. Littlejohn and W. Flynn, *Phys. Rev. A* **44**, 5239 (1991).
- [17] C. Lessig, *Modern Foundations of Light Transport Simulation*, Ph.D. thesis, University of Toronto, Toronto (2012).
- [18] G. Bal, *J. Opt. Soc. A* **23**, 1639 (2006).
- [19] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, 2nd ed. (Springer, 1989).
- [20] T. S. Ratiu and R. Schmid, *Math. Z.* **177**, 81 (1981).
- [21] M. Kline and I. W. Kay, *Electromagnetic Theory and Geometrical Optics* (John Wiley and Sons, 1965).
- [22] J. Jost, *Riemannian Geometry and Geometric Analysis*, Universitext (Springer, 2008).
- [23] D. S. Goodman, in *Handbook of Optics*, edited by M. Bass (McGraw-Hill Companies, 2010) 3rd ed., Chap. 1, pp. 1.3–1.92.
- [24] D. G. Ebin and J. E. Marsden, *Ann. Math.* **92**, 102 (1970).
- [25] V. I. Arnold, *Annales de l'institut Fourier* **16**, 319 (1966).
- [26] We disregard here some technical details in the construction of $\mathfrak{g}^*$, cf. [17, Chapter 2.3.5.3].
- [27] J. E. Marsden, A. Weinstein, T. S. Ratiu, R. Schmid, and R. G. Spencer, in *Atti Acad. Sci. Torino Cl. Sci. Fis. Math. Natur.* (1982) pp. 289–340.
- [28] J. J. Duderstadt and W. R. Martin, *Transport Theory* (Wiley, New York, 1979).
- [29] A. Gershun, *J. Math. Phys.* **18**, 51 (1939).
- [30] R. W. Preisendorfer, *Radiative Transfer on Discrete Spaces* (Pergamon Press, 1965).
- [31] See [1] and references therein.
- [32] See for example [33]. In theoretical optics, various names are employed for the Wigner transform, cf. [34].
- [33] E. Wolf, *Phys. Rev. D* **13**, 869 (1976).
- [34] M. J. Bastiaans, *J. Opt. Soc. A* **3**, 1227 (1986).
- [35] Interesting work in this direction is for example [36, 37].
- [36] C. Emmrich and A. Weinstein, *Comm. Math. Phys.* **176**, 701 (1996).
- [37] H. Bursztyn and S. Waldmann, *Lett. Math. Phys.* **53**, 349 (2000).
- [38] M. I. Mishchenko, *Journal of Quantitative Spectroscopy and Radiative Transfer* **112**, 2079 (2011).
- [39] D. D. Holm, V. Putkaradze, and C. Tronci, *Proc. R. Soc. A*, 43 (2010).
- [40] M. I. Mishchenko, L. D. Travis, and A. A. Lacis, *Multiple Scattering of Light by Particles: Radiative Transfer and Coherent Backscattering* (Cambridge University Press, 2006).
- [41] M. I. Mishchenko, in *Radiative Transfer*, edited by M. P. Menguc and N. Selcuk (Begell House, New York, 2007) pp. 1–30.